

UNIVERSITY
UTRECHT



DEPARTMENT
OF
MATHEMATICS

ON A THEOREM OF TARSKI
by
Adrian Rezus

PREPRINT
NR. 227

January 1982

ON A THEOREM OF TARSKI*

by

Adrian Rezus

In the mid-twenties Alfred Tarski raised the problem of axiomatizing classical propositional logic (\mathcal{TV}_{\wedge} , for short) by a single axiom and the rules of detachment (for material implication) and substitution as primitive rules of derivation. The problem was solved, by Tarski again, in 1925, for a large class of propositional logics, as eventually announced in [43]_{AA}, Theorem 8, but no proof of the result claimed there was ever published.

The method of Tarski for finding single axioms was frequently referred to in print (see, e.g., [9]_{AA}, [23]_{AA}, [11]_{AA}, [14]_{AA}, etc.) and though several persons — among which authors of textbooks of logic — have been certainly familiar to the principle involved in the original proof, none of them has ever documented the method for a larger audience.

We have discovered incidentally an analogue of Tarski's original method in the late 1979, relying on an extremely simple lambda-calculus argument (cf. [19]_{AA}). This note reports some elaborate details of work contained, in essence, in [19]_{AA}, section 4, extending Tarski's result in various directions.

* This work was partly supported by the University of Utrecht. The author is indebted to Henk Barendregt, David Meredith and Willem L. van der Poel for useful comments on earlier drafts of this paper. Also Ian Douglas, Robert K. Meyer (A.N.U., Canberra) and Alasdair Urquhart (University of Toronto) have largely facilitated us the access to unpublished, as yet, material on relevant logics.

1. Preliminaries.

Throughout in the sequel, a propositional language is constructed as usual, from a denumerably infinite list of propositional variables $p, q, r, s, t, u, v, w, \dots$ (possibly affected by numeric sub- and/or superscripts) and some unspecified propositional connectives. Whenever the latter are fixed we use Łukasiewicz's parentheses-free frontal notation in order to denote propositional formulae. In particular, C will be used as a binary connective and stands for any suitable notion of implication (material, intuitionistic, strict, multiple-valued, relevant, etc.).

A propositional language \mathcal{L} is implicative if some notion of implication C is either primitive in \mathcal{L} or can be defined in terms of the primitive notions in \mathcal{L} . A purely implicative language is a propositional language containing only C as a primitive (propositional) connective.

Lower-case light face Greek letters $\alpha, \beta, \gamma, \dots$ (possibly affected by sub- or superscripts) will be used as meta-variables on propositional formulae in some implicative language, modulo uniform reletterings of the propositional variables they contain.

Similarly, lower-case bold face Roman letters will be used as constants for fixed propositional formulae modulo such reletterings. For convenience, we shall pick out a standard (e.g., lexical) ordering of the propositional variables and use formulae with their propositional variables occurring in this order as ad hoc representatives for specific bold face letters.

Examples (to be used later on):

$\underline{\underline{1}} := Cpp \text{ or } := Cp_o p_o$
 $\underline{\underline{k}} := CpCqp$
 $\underline{\underline{k'}} := CpCqq$
 $\underline{\underline{b}} := CCpqCCrpCrq$
 $\underline{\underline{b'}} := CCpqCCqrCpr$
 $\underline{\underline{c}} := CCpCqrCqCpr$
 $\underline{\underline{c'}} := CpCCqCprCqr$
 $\underline{\underline{c_*}} := CpCCpqq$

$\underline{d}_0 := \underline{d} := \text{CpCqCCpCqrr}$
 $\underline{d}_n := \text{CpCqCCpCqrrCs}_1 \dots \text{Cs}_n \text{r}$
 $\underline{t} := \text{CpCqCrCCpCqCrss}$
 $\underline{s} := \text{CCpCqrCCpCqCpr}$
 $\underline{s}' := \text{CCpCqCCpCqCpr}$
 $\underline{w} := \text{CCpCpCqCpCq.}$

As a shorthand, a constant formula (denoted by/represented by some bold face letter) β may occur as a subformula of a propositional formula α . In such cases the notational convention is that no propositional variable occurring in β should occur elsewhere in α . E.g., \underline{k}^+ stands for Cpk , i.e., for CpCqCrq or some lexical variant of it, but definitely not for CpCpCrp , CpCqCpCq , etc. This will be sometimes made explicit in current auxiliary notation. Thus where

$\underline{i}_0 := \underline{i} := \text{Cp}_0 \text{p}_0, \quad \underline{i}_n := \text{Cp}_n \text{p}_n \quad (n \geq 1),$
 we will also set

$\underline{k}_0 := \underline{c}_* := \text{CpCCpqq}, \quad \underline{k}_n := \text{CpCC}\underline{i}_1 \dots \underline{i}_n \text{CCpqq} \quad (n \geq 1),$
 $\underline{k}'_0 := \text{CCCpqq}, \quad \underline{k}'_n := \text{CC}\underline{i}_1 \dots \underline{i}_n \underline{i}_{n+1} \text{qq} \quad (n \geq 1)$

and

$\underline{k}_0^+ := \text{CC}\underline{k}_0 \text{rr} := \text{CCCPCCpqqrr}, \quad \underline{k}_n^+ := \text{CC}\underline{i}_1 \dots \underline{i}_n \underline{k}_n \text{rr} \quad (n \geq 1).$

A (system of) propositional logic will be often confused with the set of its theorems, but whenever not otherwise specified, we shall understand by "propositional logic" a Hilbert-style presentation of some concept of logical derivation.

Where \underline{L} is a propositional logic, the set of its (well formed) formulae will be denoted by $\text{Form}_{\underline{L}}$.

A propositional logic \underline{L} is implicative (relative to some specified notion of implication C) if (i) so is its underlying language and, moreover, (ii) the rule of detachment for C (modus ponens; (MP), for short)

$$\text{C}\alpha\beta, \alpha \implies \beta$$

is a derivable rule in \underline{L} (say not only admissible in \underline{L} ; for the distinction derivable/admissible in \underline{L} see, e.g., [40]). A propositional logic is purely implicative if it is implicative and its underlying language is purely implicative. (This is mere technical jargon not intended to commit ourselves to some particular assumptions — philosophical or so — concerning what is to be meant by "implication" at all. But see [22], Chapter 4, for details.)

In particular, if some implicative logic is denoted by L_{λ} then its pure(ly implicative) fragment (relative to the specified notion of implication) will be denoted by $L_{\lambda} \rightarrow$.

In view of a remark of John von Neumann, it is immaterial if we choose to present a propositional logic with axioms and the rule of substitution (henceforth: (SB)) as a primitive rule of derivation or use axiom schemes and give up the rule (SB). So it will be convenient to forget any explicit reference to the applications of (SB), save in critical cases or in examples when we shall adopt the usual Polish school notation for substitutions (see, e.g., [12] or [18]).

While indicating proofs by (MP) (and (SB)) from particular sets of implicative formulae we shall make heavy use of (a slight refinement of) C.A. Meredith's condensed detachment operator (cdo, for short).

Initially thought of as a simple and convenient notational expedient (cf. [18]: Appendix II, [14], [8], [15], [16], [17], etc.), this abbreviative device has also some deeper motivation and applications as it will be seen later on.

Roughly speaking, where α, β are (purely) implicative formulae, $D_{\lambda} \alpha \beta$ stands for the most general result of the detachment of β or some substitution instance of it (as a minor premiss of (MP)) from α or some substitution instance of it (as a major premiss of (MP)). So $D_{\lambda} \alpha \beta$ makes sense for any two implicative formulae α, β such that there are substitution instances α', β' of α, β resp. with $\alpha' = C\beta'\gamma$ for some implicative formula γ . If this is the case we will say that $D_{\lambda} \alpha \beta$ is a proof of γ or that $D_{\lambda} \alpha \beta$ proves γ and it is easy to see that γ is uniquely determined up to uniform reletterings of its propositional variables. Here "the most general result" must be understood à la C.A. Meredith, in the sense that we should not make unnecessary identifications of propositional variables while performing the underlying (condensed) detachment.

Obviously, C.A. Meredith's D -meta-notation for proofs by (MP) (and (SB)) allows a non-ambiguous restoring of the missing substitutions (= applications of (SB)), modulo uniform reletterings of propositional variables.

Examples: where $\underline{i}, \underline{k}, \underline{k}', \underline{c}$ and \underline{c}_* are as earlier we have that

$\underline{D}_{\underline{c}} \underline{k}$ proves \underline{k}' ,

$\underline{D}_{\underline{c}} \underline{k} \underline{i}$ proves \underline{k}'

and

$\underline{D}_{\underline{c}} \underline{i}$ proves \underline{c}_* ,

while, with applications of (SB) written up in full, the latter proof might have been displayed in the spirit of the Polish school as follows:

```

1  :=  c      :=  CCpCqrCqCpr
2  :=  i      :=  Cpp
      1[p/Cpq, q/p, r/q] * C2[p/Cpq] - 3
3  :=  c*     :=  CCpCCppqq.

```

The stipulation requiring the most general result of a detachment forbids taking say CpCpp for the result "proved by" $\underline{D}_{\underline{c}} \underline{k}$ or $\underline{D}_{\underline{c}} \underline{i}$.

As a further notational convention we shall write $\underline{D}_{\underline{c}} \alpha \beta = \underline{D}_{\underline{c}} \alpha' \beta'$ if $\underline{D}_{\underline{c}} \alpha \beta$ and $\underline{D}_{\underline{c}} \alpha' \beta'$ prove the same formula γ (modulo the due reletterings). So one should have $\underline{D}_{\underline{c}} \underline{k} = \underline{D}_{\underline{c}} \underline{i}$.

More accurately, the cdo \underline{D} may be thought of as being a partial binary operator on sets of implicative formulae whether pure or not. We shall give here a closer description of C.A. Meredith's \underline{D} using the unification algorithm of J.A. Robinson (cf., e.g., [20], [24]). For convenience, let us restrict the frame of reference to purely implicative languages (the extension to arbitrary implicative languages is trivial).

Let \underline{L} be a set of purely implicative formulae. Then the cdo is a partial mapping

$$\underline{D}: \underline{L} \times \underline{L} \rightarrow \underline{L}$$

such that, for all α, β in \underline{L} , $\underline{D}_{\underline{c}} \alpha \beta := \underline{D}(\alpha, \beta)$ is defined if

(i) $\alpha = C \gamma' \gamma''$ and

(ii) β and the antecedent γ' of α have a unifier in the sense of [20],

else $\underline{D}_{\underline{c}} \alpha \beta$ is undefined.

If $\underline{D}_{\underline{c}} \alpha \beta$ is defined then β and the antecedent of α have a most general unifier in the sense of [20] (mgu, for short), β' say, and the due substitution instances α', β' of α, β resp. give $\alpha' = C \beta' \gamma$, for some γ , which is unique (up to uniform reletterings). Then $\underline{D}_{\underline{c}} \alpha \beta := \gamma$.

It should be noted that the definition suggested above is constructive in the sense that the mgu of every two formulae (if any) can be found effectively by the so-called unification algorithm. (Actually, the algorithm allows to establish whether or not the unification is possible and if this is the case it finds out the due mgu. For details, see [20], [21] or consult [15] for an alternative equivalent account.)

2. Meredith Δ -proofs.

For the purposes of this paper it will be convenient to formalize the meta-language of Δ -proofs and to introduce some systematic abbreviations.

For any implicative logic L , let the set of Δ -proof expressions (pe's, for short) of L be the least set D_L such that

- (i) any propositional meta-variable ($\alpha, \beta, \gamma, \dots$) is in D_L ,
- (ii) any propositional constant (i.e., a bold face letter denoting a theorem of L) is in D_L ,
- (iii) if x, y are in D_L then so is Δxy .

Hereafter, x, y, z, \dots (possibly with sub- or superscripts) will range on pe's of some arbitrary implicative logic L .

A pe consisting of a single letter (propositional meta-variable or propositional constant) is atomic.

Further we write (xy) for Δxy and save parentheses by omitting the outermost pair and assuming association to the left.

E.g., $\underline{\underline{d}}(\underline{\underline{k}}\underline{\underline{k}})(\underline{\underline{d}}\underline{\underline{k}}\underline{\underline{i}})$ stands for $\underline{\underline{\Delta\Delta d\Delta k\Delta k\Delta d\Delta k i}}$.

We also adopt the following use of numeric superscripts ($n \geq 0$): for any atomic pe x of some implicative logic L ,

- x^0 is the empty word,
- $x^1 \equiv x$ and $x^{n+1} \equiv x^n x$, for all $n \geq 0$.

Similarly, we write C^n ($n \geq 0$) for n consecutive occurrences of C in some implicative formula, $(CC)^n$ for $2 \times n$ consecutive occurrences of C , etc. But p_j^i, q_j^i, \dots ($i, j \geq 0$) are propositional variables and the superscript "i" has no iterative effect.


If a pe x is defined ("meaningful", "denoting") we write x^\downarrow to indicate this, otherwise (if x is not defined) we write x^\uparrow . It is reasonable to assume that a pe x is defined whenever it is atomic, so we won't write $\alpha^\downarrow, \beta^\downarrow$, etc. and $\underline{\underline{k}}^\downarrow, \underline{\underline{i}}^\downarrow$, etc. either.

Examples: $\underline{\underline{s}}\underline{\underline{i}}\underline{\underline{i}}^\uparrow, \underline{\underline{w}}\underline{\underline{i}}^\uparrow$, but $\underline{\underline{i}}x^\downarrow, \underline{\underline{k}}x^\downarrow, \underline{\underline{k}}'x^\downarrow$ whenever x^\downarrow (if x is atomic say).

In evaluating pe's we may adopt some obvious inside-out and left-to-right strategy which can be always represented in a tree-like manner.

E.g., with the example above: $\underline{d}(\underline{kk})(\underline{dki})$, we should first evaluate $\underline{kk}, \underline{dk}$, next $\underline{d}(\underline{kk})$, then \underline{dki} and, finally, the entire pe.

Tree-like picture:



It should be clear how to define components of pe's (or sub- λ -proof expressions; sub-pe's, for short). Further, a proper component (sub-pe) of a pe x is any component of x distinct from x itself. (Sub-pe's of a pe correspond, as expected, to subtrees, in the tree-representation of a pe.)

Now the evaluation of a pe depends on the evaluation of its components and it should respect in some immediate sense a variant of the Frege principle of significance. For, obviously, if some component x of a pe y is such that x^\uparrow then the entire pe y is such that y^\uparrow , while if every proper component x of y is such that x^\downarrow then one should also have y^\downarrow (otherwise, x is a component of x , for any pe x).

So far we have introduced a class of (interpreted) formal languages suited for some slightly modified typed combinatory logics (cf. [4], [6] or [2]: Appendices A, B). To see how they work we need not develop special theories of reduction for each implicative logic λ but consider only the process of evaluation of pe's.

For any two pe's x, y define $x = y$ if $x^\downarrow, y^\downarrow$ and x and y have the same value; clearly, $=$ is an equivalence relation on pe's z with z^\downarrow . (So, to make a parallel with the classical case, the evaluation of a pe may be compared with a kind of "combinatory reduction", while $=$ is supposed to be introduced by the familiar Church-Rosser property, where " $x = y$ " means that x and y have a common "reduct". In fact, our pe's have the so-called "strong normalization property" whenever they are defined, i.e., make sense as "typed terms", and hence

our choice of "evaluation" — and not "reduction" — as a central concept in the "theory of D_{\wedge} -proofs". See also [4], 9E and [7].)

Now $=$ can be extended in the obvious way up to a partial equivalence relation \simeq on pe's, and we will want to write " $x \simeq y$ " for any two pe's x, y , whether they are defined ("meaningful") or not. (For later reference note that we have introduced \simeq starting from the definition of $=$ and not conversely.)

The relations $=$ and \simeq may be interpreted intuitively as equivalence (partial equivalence) of proofs (by (MP) and (SB)) in some implicative logic L_{\wedge} .

As earlier, if x, y are pe's of some L_{\wedge} and y is atomic then y^{\downarrow} and we say that x proves y whenever $x = y$. Also note that, by construction, definiteness of pe's (\dots^{\downarrow}) is preserved by $=$. So for all pe's x, y if x proves y then also x^{\downarrow} (for y is atomic, hence y^{\downarrow} , while $=$ preserves definiteness).

It is easy to check the following combinatory-like "equations":

LEMMA 1.

For all pe's x, y, z, z_1, \dots, z_n ($n \geq 1$):

- | | |
|---|--|
| (1) $\underline{i}x \simeq x$ | (2) $\underline{k}xy \simeq x$ |
| (3) $\underline{k}'xy \simeq y$ | (4) $\underline{k}^+xyz \simeq y$ |
| (5) $\underline{b}xyz \simeq x(yz)$ | (6) $\underline{b}'xyz \simeq y(xz)$ |
| (7) $\underline{c}xyz \simeq xzy$ | (8) $\underline{c}'xyz \simeq yzx$ |
| (9) $\underline{c}_*xy \simeq yx$ | (10) $\underline{d}xyz \simeq zxy$ |
| (11) $\underline{d}_nxyzz_1 \dots z_n \simeq zxy$ | |
| (12) $\underline{t}xyz \simeq xzy$ | (13) $\underline{w}xy \simeq xyy$ |
| (14) $\underline{s}xyz \simeq xz(yz)$ | (14) $\underline{s}'xyz \simeq yz(xz)$. |

Proof. Straightforward consequences from the definition of D_{\wedge} . \square

Now $=$ and \simeq are congruences w.r.t. D_{\wedge} . That is:

LEMMA 2.

For all pe's x, y, z ,

- (1) if $x = y$ and xz^{\downarrow} (or yz^{\downarrow}) then $xz = yz$,
 - (2) if $x = y$ and zx^{\downarrow} (or zy^{\downarrow}) then $zx = zy$,
- and, finally,
- (3) if $x \simeq y$ then $xz \simeq yz$ and $zx \simeq zy$.

Proof. Obvious. \square

REMARK 3.

For all $n \geq 0$,

- (1) if x^\downarrow and y^\downarrow then $\underline{d}_n x^\downarrow$ and $\underline{d}_n xy^\downarrow$,
- (2) if $x^\downarrow, y^\downarrow$ and z^\downarrow then $\underline{t}x^\downarrow, \underline{t}xy^\downarrow$ and $\underline{t}xyz^\downarrow$.

Some of the following consequences from definitions will be useful later on.

LEMMA 4.

For all implicative formulae α, β in some L_A and all $n \geq 0$,

- (1) $\underline{i}\alpha = \alpha$
- (2) $\underline{k}\alpha\beta = \alpha$
- (3) $\underline{k}'\alpha\beta = \beta$
- (4) $\underline{k}\underline{i} = \underline{c}\underline{k} = \underline{k}'$
- (5) $\underline{k}\underline{i}\alpha = \underline{c}\underline{k}\alpha = \underline{k}'\alpha = \underline{i}$
- (6) $\underline{k}\underline{k} = \underline{k}^+$
- (7) $\underline{k}_n\alpha\beta = \beta \underline{i}^n \alpha$
- (8) $\underline{k}'_n\alpha = \alpha \underline{i}^{n+1}$
- (9) $\underline{k}_n^+\alpha = \alpha \underline{i}^n \underline{k}_n$

Proof. Straightforward. \square

LEMMA 5.

- (1) $\underline{c}\underline{b} = \underline{b}'$
- (2) $\underline{c}\underline{b}' = \underline{b}'\underline{c}_*(\underline{b}'\underline{b}') = \underline{b}$
- (3) $\underline{b}'\underline{b}'(\underline{b}'\underline{c}_*) = \underline{b}(\underline{c}_*\underline{c}')\underline{c}' = \underline{c}'\underline{c}'\underline{c}' = \underline{c}$
- (4) $\underline{b}\underline{b}\underline{c}_* = \underline{c}\underline{c} = \underline{c}'$
- (5) $\underline{c}\underline{i} = \underline{c}_*$
- (6) $\underline{c}\underline{s} = \underline{s}'$
- (7) $\underline{c}\underline{s}' = \underline{s}$.

Proof. Easy. \square

LEMMA 6.

For all $n \geq 0, m \geq 2$,

- (1) $\underline{k}_n \underline{i} = \underline{k}'_n$
- (2) $\underline{k}_n \underline{k}_n = \underline{k}_n^+$
- (3) $\underline{k}_n \underline{i} \underline{i} = \underline{i}$
- (4) $\underline{k}_n \underline{i}^m = \underline{i}$.

Proof. Easy. \square

Where x, y are pe's, let $x[[y]]$ stand for " y is a component of x ". Then we have a

COROLLARY 7.

If $x[[y]] \simeq z$ and $y \simeq y_0$ then $x[[y_0]] \simeq z$, for all pe's x, y, y_0, z .

Proof. Use Lemma 2. \square

The evaluation strategy indicated earlier is guaranteed by the following consequence of Corollary 7.

COROLLARY 8.

For all pe's x, y, y_0, z of some L_λ ,
 if $x[y]$ proves z and y proves y_0 then $x[y_0]$ proves (also) z .
Proof. Indeed, if $x[y]$ proves z then z is atomic and one has z^\downarrow , by definition. So $x[y]^\downarrow$ and also y^\downarrow , by the "Frege principle" noted earlier; hence y has a "value". But y proves y_0 so this "value" must be y_0 . So $x[y]$ and $x[y_0]$ must have the same "value", by Corollary 7. \square

Let L_λ be some implicative logic, If β is a theorem of L_λ and

$$\beta_{\underline{i}}^m = \underline{i}$$

for some $m \geq 0$, we say that β is m-solvable. Similarly, a set $\{\beta_j : j \in I\}$ ($I \subseteq \mathbb{N}$) of theorems of L_λ is m-solvable if

$$\beta_{j\underline{i}}^m = \underline{i}$$

for all j in $I \subseteq \mathbb{N}$.

Then one has immediately

LEMMA 9.

For all α, β in some implicative logic L_λ ,

- (1) if β is m-solvable ($m > 0$) then $k_m \alpha \beta = \alpha$,
- (2) k_m is m-solvable, for $m \geq 2$,
- (3) $k_m \alpha k_m = \alpha$, for all $m \geq 2$.

Proof. (1) Use Lemma 4: (6) and (1) with Corollary 8.

(2) Note that $k_m \underline{i} = \underline{i}^m \underline{i} = \underline{i}$, for all $m \geq 0$.

(3) By (1) and (2). \square

Let L_λ be an arbitrary implicative logic. Define, for each $m, n \geq 1$, mappings

$$\begin{aligned} d_{m,n} : \text{Form}_{L_\lambda}^n &\longrightarrow \text{Form}_{L_\lambda}, \\ \text{by: for all } \beta_1, \dots, \beta_n &\text{ in } \text{Form}_{L_\lambda}, \text{ not containing } p_j^i, q_j^i \text{ (} 1 \leq i \leq n-1, \\ &1 \leq j \leq m), \\ d_{m,n}(\beta_1, \dots, \beta_n) &:= \begin{cases} (CC)^{n-1} \beta_1 C \beta_2 p_1 C q_1^1 \dots C q_m^1 p_1 \dots C \beta_n p_{n-1} C q_1^{n-1} \dots C q_m^{n-1} p_{n-1} & \text{if } n > 1, \\ \beta_1, & \text{if } n = 1. \end{cases} \end{aligned}$$

In particular, set for $m = 0$, $d_n := d_{0,n}$, with, for all β_1, \dots, β_n in Form_{L_λ} , not containing p_i ($1 \leq i \leq n-1$),

$$\tilde{d}_n(\beta_1, \dots, \beta_n) := \begin{cases} (CC)^{n-1} \beta_1 C \beta_2 p_1 p_1 \dots C \beta_n p_{n-1} p_{n-1}, & \text{if } n > 1 \\ \beta_1, & \text{if } n = 1. \end{cases}$$

LEMMA 10.

(1) For all $\alpha_1, \dots, \alpha_n$ in some L_λ ($n \geq 1$), not containing p_1, \dots, p_{n-1} , one has

$$\tilde{d}_m^{n-1} \alpha_1 \dots \alpha_n \text{ proves } \tilde{d}_n(\alpha_1, \dots, \alpha_n).$$

(2) For all $\alpha_1, \dots, \alpha_n$ in some L_λ and all $m > 0, n \geq 1$, where the α_k 's do not contain p_j^i, q_j^i ($1 \leq i \leq n-1, 1 \leq j \leq m$), and where we have set $\tilde{d}_m^{n-1} := (\tilde{d}_m)^{n-1}$,

$$\tilde{d}_m^{n-1} \alpha_1 \dots \alpha_n \text{ proves } \tilde{d}_{m,n}(\alpha_1, \dots, \alpha_n).$$

Proof. Tedious but trivial. \square

Let also L_λ be as earlier and t_λ be a mapping

$$t_\lambda : \text{Form}_{L_\lambda}^3 \longrightarrow \text{Form}_{L_\lambda}$$

such that, for all $\beta_1, \beta_2, \beta_3$ in Form_{L_λ} not containing p ,

$$t_\lambda(\beta_1, \beta_2, \beta_3) := CC\beta_1 C\beta_2 C\beta_3 pp.$$

Then one can see easily that

LEMMA 11.

For all $\alpha_1, \alpha_2, \alpha_3$ in some implicative logic L_λ ,

$$\tilde{t}_{\alpha_1 \alpha_2 \alpha_3} \text{ proves } t_\lambda(\alpha_1, \alpha_2, \alpha_3).$$

Proof. Clear. \square

3.A generalization of Tarski's theorem.

Let L_{λ} be a propositional logic. Where $(R_1), \dots, (R_r)$, $(r \geq 1)$ are derivable rules in L_{λ} we say that L_{λ} is finitely axiomatizable in the set of rules $\{(R_1), \dots, (R_r)\}$ if L_{λ} has a Hilbert-style formulation with

- (i) a finite number of axioms $\alpha_1, \dots, \alpha_n$, $(n \geq 1)$, and
- (ii) the rules $(R_1), \dots, (R_r)$, (SB) as primitive rules of derivation.

Let now L_{λ} be an implicative logic with (MP), $(R_1), \dots, (R_r)$, $(r \geq 0)$ derivable rules in L_{λ} . We say that L_{λ} is Tarski axiomatizable in the set of rules $\{(R_1), \dots, (R_r), (MP)\}$ if L_{λ} has a Hilbert-style formulation with

- (i) a single axiom and
- (ii) (MP), for the specified notion of implication, together with
- (iii) $(R_1), \dots, (R_r)$, (SB) as primitive rules of derivation.

(Alternatively, one can take a single/a finite number of axiom scheme/s and leave out the rule (SB), as indicated earlier.)

Clearly, for L_{λ} implicative, if L_{λ} is Tarski axiomatizable in some set of rules then it is also finitely axiomatizable in the same set. We establish sufficient conditions, generalizing Theorem 8 in [13], and allowing to prove the converse of the above.

THEOREM 12.

Let L_{λ} be an implicative logic (relative to some specified notion C of implication). If

- (i) L_{λ} is finitely axiomatizable in the set of rules $\{(R_1), \dots, (R_r), (MP)\}$ ($r \geq 0$)

and

- (ii) for some $m \geq 0$, d_m is a theorem of L_{λ} ,
 - (iii) k is a theorem of L_{λ}
- then L_{λ} is Tarski axiomatizable in the same set of rules.

Proof. Let $L_{\lambda n}$ be the (Hilbert-style) formulation of L_{λ} with axioms $\alpha_1, \dots, \alpha_n$ ($n \geq 1$)

and rules

$$(R_1), \dots, (R_r), (MP), (SB) \quad (r \geq 0).$$

(If $n = 1$, there is nothing to prove but we include this as a limit case.)

Define, for $m \geq 0$ and $n \geq 1$, (m, n fixed),

$$\underline{h}_{m,n} := \underline{d}_{m,n}(\alpha_1, \dots, \alpha_n).$$

Then, by Lemma 10, one finds that

$$\underline{d}_m^{n-1} \alpha_1 \dots \alpha_n \text{ proves } \underline{h}_{m,n}$$

and since \underline{d}_m is a theorem of \underline{L}_n , $\underline{h}_{m,n}$ is also a theorem of \underline{L}_n (for $\underline{h}_{m,n}$ can be proved by (MP) and (SB) from the axioms of \underline{L}_n ; hence the result holds for $r = 0$, too).

Now set

$$\underline{g}_{m,n} := \underline{d}_{m,3}(\underline{k}, \underline{k}^+, \underline{h}_{m,n}),$$

with $\underline{k}, \underline{k}^+$ and $\underline{h}_{m,n}$ as above.

Our claim is that $\underline{g}_{m,n}$ is the needed single axiom. Firstly, $\underline{g}_{m,n}$ is a theorem of \underline{L}_n . Indeed, by Lemma 4:(6), one finds that

$$\underline{k}\underline{k} \text{ proves } \underline{k}^+$$

and, by Lemma 10,

$$\underline{d}_m \underline{d}_m \underline{k}\underline{k}^+ \underline{h}_{m,n} \text{ proves } \underline{g}_{m,n},$$

hence the result follows by Corollary 8 (for $\underline{k}, \underline{d}_m$ are theorems of \underline{L}_n , by hypothesis).

On the other hand, let \underline{L}_n^* be the (Hilbert-style) formulation of \underline{L}_n with $\underline{g}_{m,n}$ as single axiom and the same rules as earlier.

(Explicitly, $\underline{g}_{m,n}$ is

$$\underline{g}_{m,n} := \text{CCCC}\underline{k}\underline{k}^+ \underline{r}_1 \underline{C} \underline{s}_1 \dots \underline{C} \underline{s}_m \underline{r}_1 \underline{C} \underline{h}_{m,n} \underline{r}_2 \underline{C} \underline{t}_1 \dots \underline{C} \underline{t}_m \underline{r}_2.)$$

Using \underline{D} -proofs one can show quickly that

$$\underline{g}_{m,n} \underline{g}_{m,n} (\underline{g}_{m,n})^m \text{ proves } \underline{k},$$

and, in particular, for $m = 0$ and with $\underline{g}_n := \underline{g}_{0,n}$, for convenience, one finds that

$$\underline{g}_n \underline{g}_n \text{ proves } \underline{k}.$$

(It is a tedious affair to find explicitly the needed substitutions, but the matter is completely trivial and we have only to take some care in applying correctly J.A. Robinson's unification algorithm. For instance, displaying substitutions à la Łukasiewicz [12], one has, with $\underline{k} := \underline{C} \underline{p}_1 \underline{C} \underline{q}_1 \underline{p}_1$ and $\underline{h}_n := \underline{h}_{0,n}$,

$$1 := \underline{g}_n := \text{CCCC} \underline{p} \underline{C} \underline{q} \underline{p} \underline{C} \underline{r} \underline{C} \underline{s} \underline{C} \underline{t} \underline{s} \underline{u} \underline{u} \underline{C} \underline{h}_n \underline{v} \underline{v}$$

$$2 := \underline{g}'_n := 1 [\underline{p} / \underline{C} \underline{h}_n \underline{k} \underline{C} \underline{h}_n \underline{C} \underline{h}_n \underline{k}, \underline{q} / \underline{C} \underline{r} \underline{C} \underline{s} \underline{C} \underline{t} \underline{s}, \underline{r} / \underline{h}_n, \underline{s} / \underline{p}_1, \underline{t} / \underline{q}_1, \\ \underline{u} / \underline{C} \underline{h}_n \underline{C} \underline{h}_n \underline{k}, \underline{v} / \underline{k}]$$

$$3 := \underline{g}''_n := 1 [\underline{p} / \underline{C} \underline{h}_n \underline{k}, \underline{q} / \underline{q}, \underline{r} / \underline{r}, \underline{s} / \underline{s}, \underline{t} / \underline{t}, \underline{u} / \underline{C} \underline{h}_n \underline{k} \underline{C} \underline{h}_n \underline{C} \underline{h}_n \underline{k}, \underline{v} / \underline{C} \underline{h}_n \underline{k}] \\ 2 * 3 = 4$$

$$4 := \underline{k} := \underline{C} \underline{p}_1 \underline{C} \underline{q}_1 \underline{p}_1,$$

$$\text{i.e., } \underline{g}'_n := \underline{C} \underline{g}''_n \underline{k}.)$$

Now, $\underline{h}_{m,n}$ is a theorem of \underline{L}_Λ^* for one can establish that $\underline{g}_{m,n}(\underline{kk})_{\underline{k}}^{m+1}$ proves $\underline{h}_{m,n}$, while the axioms $\alpha_1, \dots, \alpha_n$ of $\underline{L}_{\Lambda n}$ can be extracted from $\underline{h}_{m,n}$ as follows.

With, for $1 \leq j \leq n-1$, set

$$\underline{h}_{m,n-j} := \underline{h}_{m,n} \underline{k}^{(m+1) \times j}$$

(that is:

$$\begin{aligned} \underline{h}_{m,n-1} &:= \underline{h}_{m,n} \underline{k} \underline{k}^m = \underline{h}_{m,n} \underline{k}^{m+1} \\ \underline{h}_{m,n-2} &:= \underline{h}_{m,n-1} \underline{k} \underline{k}^m = \underline{h}_{m,n} \underline{k}^{(m+1) \times 2} \\ \dots &\dots \dots \dots \dots \dots \\ \underline{h}_{m,1} &:= \underline{h}_{m,2} \underline{k} \underline{k}^m = \underline{h}_{m,n} \underline{k}^{(m+1) \times (n-1)} \end{aligned}$$

It is easy to see that, for $1 \leq j \leq n$,

$$\underline{h}_{m,j}(\underline{kk})_{\underline{k}}^{m+1} \text{ proves } \alpha_j,$$

so the axioms of $\underline{L}_{\Lambda n}$ can be proved from $\underline{g}_{m,n}$ by (MP) and (SB) only (i.e., for $r = 0$, too).

(These axioms imply also that \underline{d}_m is a theorem of \underline{L}_Λ^* , by hypothesis.)

So $\underline{L}_{\Lambda n}$ and \underline{L}_Λ^* are equivalent. \square

REMARK 13.

Tarski's Theorem 8 in [13] is a particular case of our Theorem 12 with $m = 0$ and $n = 1$.

REMARK 14.

Our method of proving Theorem 12 does not provide organic axioms, in the sense of M. Wajsberg (for an axiom system \underline{L}_Λ , an axiom of \underline{L}_Λ is organic if it has no subformulae, except itself, that are theorems in \underline{L}_Λ) and this was also the case with Tarski's original method of proof, as reported in [11]. (The import of an organic axiomatization is explained in [24].)

Another practical inconvenient of both methods (see [23] and [11] for Tarski's examples) is in the fact the single axioms obtained thereby are very long.

In the end, Theorem 12 is of some theoretical interest since there are systems of propositional logic that are finitely axiomatizable in (MP) and still not Tarski axiomatizable in (MP). E.g., the purely implicative fragment $\underline{T}_\rightarrow$ of the logic of "Ticket Entailment" of A.R. Anderson (cf. [1]) cannot be axiomatized with a single axiom, (MP) and (SB) only. (This result is due to Z. Parks; see [1], 8.5.2., for details.)

REMARK 15.

If we had considered the additional condition

(iv) \underline{t} is a theorem of \underline{L}

among the hypotheses of Theorem 12, the construction of the single axiom $\underline{g}_{m,n}$ might have been somewhat different.

Indeed, with $\underline{h}_{m,n}$ as earlier, set

$$\underline{g}_{m,n}^* := \underline{t}(\underline{k}, \underline{h}_{m,n}, \underline{k}) := \underline{CCKCh}_{m,n} \underline{CKrr}$$

for the new single axiom.

Now, as

$$\underline{tkh}_{m,n} \underline{k} \text{ proves } \underline{g}_{m,n}^*$$

by Lemma 11, $\underline{g}_{m,n}^*$ is also a theorem of (the new) \underline{L} .

Let \underline{L}^* be the new formulation of \underline{L} with $\underline{g}_{m,n}^*$ as single axiom and primitive rules as earlier.

It is easy to see that

$$\underline{g}_{m,n}^* \underline{g}_{m,n}^* \underline{g}_{m,n}^* \text{ proves } \underline{k}$$

and

$$\underline{g}_{m,n}^* (\underline{g}_{m,n}^* \underline{g}_{m,n}^*) \text{ proves } \underline{h}_{m,n}$$

(for this note that

$$\underline{g}_{m,n}^* \underline{g}_{m,n}^* = \underline{kk} = \underline{k}^+),$$

while the α_j 's ($1 \leq i \leq n$) can be obtained from $\underline{h}_{m,n}$, with \underline{k} readily available, as in the proof of Theorem 12.

(The combinatory argument behind the construction of $\underline{g}_{m,n}^*$ is due, in essence, to J.B. Rosser [unpublished]; cf. [19], 1.4. and Theorem 19 below.)

Let $\underline{BCK}_{\rightarrow}$ be the Meredith (purely) implicative logic (cf. [18], Appendix I, [14], [17], etc.), formulated with (MP), (SB) as primitive rules and axioms $\underline{b}, \underline{c}, \underline{k}$ (for alternative axiomatizations see [14] and [19]). As pointed out by H.B. Curry, K. Iséki, R. Routley, R.K. Meyer, D. Meredith et al. there is some interest in studying $\underline{BCK}_{\rightarrow}$ and its extensions from both a combinatory and an algebraic point of view (cf. the references in [19]). But it is also a logical landmark in axiomatization problems. Indeed, one has the following consequence of Theorem 12 above.

COROLLARY 16.

Any finitely axiomatizable extension of $\underline{BCK}_{\rightarrow}$ in some set of rules $\{(R_1), \dots, (R_r), (MP)\}$, ($r \geq 0$), is Tarski axiomatizable in the same set of rules.

Proof. Note that

$\underline{\underline{ckk}}$ proves \underline{i} , Lemma 4:(5)

$\underline{\underline{ci}}$ proves $\underline{c_*}$, Lemma 5:(5)

$\underline{\underline{bcc_*}}$ proves $\underline{d_0} := \underline{d}$,

then apply Theorem 12, with $m = 0$.

Alternatively, one has also that

$\underline{b(\underline{bc})(\underline{\underline{bcc_*}})}$ proves \underline{t} ,

so \underline{t} is a theorem of $\text{BCK}_{\wedge\wedge\rightarrow}$ and one can apply the argument of Remark 15 above. \square

Now $\text{BCK}_{\wedge\wedge\rightarrow}$ is known to be a subsystem of many familiar (propositional) logics. The following list is far of being complete.

COROLLARY 17.

The following logics are Tarski axiomatizable in (MP):

- (i) the classical logic $\text{TV}_{\wedge\wedge\rightarrow}$ and its purely implicative fragment $\text{TV}_{\wedge\wedge\rightarrow}$ (cf. [13], [11], [18], etc.);
- (ii) the intuitionistic logic $\text{H}_{\wedge\wedge\rightarrow}$ and its purely implicative fragment $\text{H}_{\wedge\wedge\rightarrow}$ (cf. [5]);
- (iii) Hilbert's positive logic, Johansson's Minimalkalkül and any (finitely axiomatizable) intermediate logic (in (MP); see [5]),
- (iv) Łukasiewicz's many-valued logics Ł_n ($n \geq 3$) and Ł_{\aleph_0} (cf. [13]), etc.

Proof. Trivial derivations, using Corollary 16. \square

REMARK 18.

Arguments similar to those used earlier apply, mutatis mutandis, to quantificational extensions of the logics named above. (Do the exercises of [19], section 4.)

4. Refinements for relevant logics.

A.N. Prior noticed (cf. [14], page 181) that the original methods of Tarski for obtaining single axioms do not work in the absence of the "paradoxical" Law of Simplification \underline{k} ($:= \text{CpCqp}$). In particular, this comment applies to several interesting (implicative) logics among which the relevant logics R_{\wedge} , E_{\wedge} and some of their neighbours or rivals (see [1], [22], [28], etc.).

It will be clear from what follows that Prior's statement no longer holds for our methods.

Actually, the problem of axiomatizing Church's weak implication (in [3], i.e., the system R_{\wedge} of [1]) was raised incidentally in [1], 8.5.1. In [19] we claimed that R_{\wedge} and the Anderson-Belnap Pure Entailment system E_{\wedge} of [1] are Tarski axiomatizable in (MP) but the effective example of single axiom suggested there, for R_{\wedge} , contained an oversight.

In this section we shall state explicitly — this was not the case in [19] — some alternative lists of conditions guaranteeing the Tarski axiomatizability of a large class of (purely implicative) relevant logics (in the sense of [1], [22]), among which R_{\wedge} , E_{\wedge} etc.

We shall first introduce some convenient terminology.

Let L_{\wedge} be a (purely) implicative logic. A theorem of L_{\wedge} is solvable if it is m -solvable for some $m \geq 0$, otherwise it is unsolvable. Sets of theorems of L_{\wedge} will be referred to similarly.

Examples: $\underline{b}, \underline{b'}, \underline{c}, \underline{c'}, \underline{i}, \underline{k}, \underline{k}_m$ ($m \geq 0$) are solvable, while $\underline{w}, \underline{s}, \underline{s'}$ are unsolvable.

Clearly, unsolvable sets may contain solvable elements, but we needn't distinguish among such subtleties.

Let L_{\wedge} (implicative) be now finitely axiomatizable in some set of rules $\{(R_1), \dots, (R_r), (MP)\}$ — (MP) for the specified notion of implication — with $r \geq 0$, and all its axioms in the set

$$\underline{B}(L_{\wedge}) = \{\alpha_1, \dots, \alpha_p, \beta_1, \dots, \beta_q\}$$

($p, q \geq 0, p + q \geq 1$) such that the α_i 's ($1 \leq i \leq p$) are unsolvable and the β_j 's ($1 \leq j \leq q$) are solvable. Usually, $\underline{B}(L_{\wedge})$ is called a basis for/of L_{\wedge} . Hereafter, " $\underline{B}(L_{\wedge})$ " will refer to this description with p, q

varying as indicated ($n := p+q$ positive, with possibly either $p = 0$ or $q = 0$; so no basis is empty, but it may contain no unsolvable, resp. no solvable elements).

Consider now the functions $d_n := d_{0,n}$ of section 2. Let $m \geq 0$.

A basis $B(L)$ for some implicative logic L is sequentially m-quasi-solvable if its elements can be arranged in a sequence

$$\alpha_1, \dots, \alpha_p, \beta_1, \dots, \beta_q \quad (p+q \geq 1; p, q \geq 0),$$

with the α_j 's unsolvable, and there are theorems

$$\gamma_1, \dots, \gamma_p$$

of L such that the following formulae are m-solvable:

- (i) $\gamma_i, \beta_j, \quad (1 \leq i \leq p, 1 \leq j \leq q)$
- (ii) $\underline{v}_i := d_2(\alpha_i, \gamma_i) \quad (1 \leq i \leq p)$
- (iii) $\underline{f}_i := d_i(\underline{v}_1, \dots, \underline{v}_i) \quad (1 \leq i \leq p)$
- (iv) $\underline{h}_j := d_{j+1}(\underline{f}_p, \beta_1, \dots, \beta_j) \quad (1 \leq j \leq q).$

(Note that $\underline{f}_1 := \underline{v}_1 := d_2(\alpha_1, \gamma_1)$ and $\underline{h}_1 := d_{p+1}(\underline{v}_1, \dots, \underline{v}_p, \beta_1)$, by the definition of d_1, d_2, \dots, d_{p+1} .)

In particular, $B(L)$ is sequentially m-solvable if $p = 0$; that is, $B(L) = \{\beta_1, \dots, \beta_q\}$ contains only m-solvable axioms and for some permutation π of the set $\{1, \dots, q\}$, the formulae

$$\underline{h}_j^\pi := d_j(\beta_{\pi(1)}, \dots, \beta_{\pi(j)}) \quad (1 \leq j \leq q)$$

are also m-solvable (viz., \underline{h}_j^π proves \underline{v}_j , for all $j, 1 \leq j \leq q$).

For simplicity, we shall consider first (purely) implicative logics possessing at least one (sequentially) m-solvable basis, for some $m \geq 2$. Next we shall extend our discussion to implicative logics possessing arbitrary bases.

THEOREM 19.

Let L_λ be an implicative logic (for some specified notion C of implication) such that

- (i) L_λ is finitely axiomatizable in the set of rules
 $\{(R_1), \dots, (R_r), (MP)\}$ ($r \geq 0$),

If, for some $m \geq 2$,

- (ii) L_λ has a sequentially m -solvable basis

- (iii) k_m is a theorem of L_λ ,

and

- (iv) d, t are theorems of L_λ

then L_λ is Tarski axiomatizable in the same set.

Proof. Let $\alpha_1, \dots, \alpha_n$ ($n \geq 1$) be the axioms of L_λ (taking $n = 1$, trivially, as a limit case of the Theorem) and construct, for an appropriate π , formulae

$$h_{\pi}^{\pi} := d_j(\alpha_{\pi(1)}, \dots, \alpha_{\pi(j)}) \quad (1 \leq j \leq n).$$

Sequential m -solvability means that, for some π , one has

$$\alpha_j \text{ proves } i \quad (1 \leq j \leq n),$$

$$h_{\pi}^{\pi} \text{ proves } i \quad (1 \leq j \leq n).$$

Now recall that, by Lemma 9 : (2) we have

$$k_m \text{ proves } i, \quad \text{for all } m \geq 2.$$

We claim the needed single axiom is

$$g_{(m,n)}^{\pi} := t(k_m, h_{\pi}^{\pi}, k_m) := CCk_m Ch_{\pi}^{\pi} Ck_m pp$$

(observing the relettering convention of section 1 above).

Indeed, k_m, d are theorems of L_λ (the hypothesis of the Theorem), so h_{π}^{π} is a theorem of L_λ , for, by Lemma 10,

$$d^{n-1} \alpha_1 \dots \alpha_n \text{ proves } h_{\pi}^{\pi}, \quad \text{for any } \pi.$$

As t is a theorem of L_λ (by hypothesis, again), one has also that, for an appropriate π , $g_{(m,n)}^{\pi}$ is a theorem of L_λ , since

$$tk_m h_{\pi}^{\pi} k_m \text{ proves } g_{(m,n)}^{\pi},$$

by Lemma 11.

Conversely, let $L_{\lambda*}$ be the formulation of L_λ with $g_* := g_{(m,n)}^{\pi}$, for convenience (m, n fixed), and primitive rules as earlier.

One works in $L_{\lambda*}$, deriving first k_m from g_* , next h_{π}^{π} and, finally, the α_j 's ($1 \leq j \leq n$). This can be done as follows:

$$g_* g_* \text{ proves } k_m^+,$$

for h_{π}^{π} is m -solvable, by the hypothesis of the Theorem.

Recall also that, by Lemma 6:(2),

$$\underline{k}_m \underline{k}_m \text{ proves } \underline{k}_m^+, \quad (m \geq 0),$$

so

$$\underline{g}_* \underline{g}_* = \underline{k}_m \underline{k}_m = \underline{k}_m^+, \quad (m \geq 2)$$

and hence

$$\underline{g}_* \underline{g}_* \underline{k}_m^+ = \underline{k}_m^+ \underline{k}_m^+ = \underline{k}_m \quad (m > 2),$$

given the m -solvability of \underline{k}_m for $m \geq 2$, (Lemma 9:(2) above), while

$$\underline{g}_* \underline{g}_* \underline{k}_m^+ \text{ proves } \underline{i} \quad \text{for } m = 2,$$

and therefore

$$\underline{g}_* \underline{g}_* \underline{i} = \underline{k}_m \underline{k}_m \underline{i} = \underline{i} \underline{i}^m \underline{k}_m = \underline{k}_m \quad (m = 2),$$

by Lemma 4:(7) and Corollary 7, etc.

So far we have shown that \underline{k}_m is a theorem of L_* , for all $m \geq 2$.

Now

$$\underline{g}_* \underline{k}_m^+ = \underline{g}_* (\underline{k}_m \underline{k}_m) = \underline{h}_n^\pi,$$

for \underline{k}_m is m -solvable ($m \geq 2$), by Lemma 9:(2).

That is, collecting the facts,

$$\underline{g}_* (\underline{g}_* \underline{g}_*) \text{ proves } \underline{h}_n^\pi \quad \text{for } m \geq 2$$

so, anyway, $\underline{k}_m, \underline{k}_m^+$ and \underline{h}_n^π can be proved from \underline{g}_* by (MP) and (SB) only and the result holds for $r = 0$, too.

Next, since $\alpha_j \underline{i}^m$ proves \underline{i} , for $1 \leq j \leq m$, we have

$$\begin{aligned} \underline{h}_{n-m}^\pi \underline{k}_m &= \underline{h}_{n-1}^\pi \\ \underline{h}_{n-1}^\pi \underline{k}_m &= \underline{h}_{n-m}^{\pi 2} = \underline{h}_{n-2}^\pi \\ \dots \dots \dots \\ \underline{h}_2^\pi \underline{k}_m &= \underline{h}_{n-m}^{\pi n-1} = \underline{h}_1^\pi = \alpha_1. \end{aligned}$$

Finally, for all $j, 1 \leq j \leq n$,

$$\underline{h}_j^\pi \underline{k}_m^+ \underline{k}_m \text{ proves } \alpha_j \quad (m \geq 2),$$

since

$$\underline{h}_j^\pi \underline{i}^m \text{ proves } \underline{i} \quad (1 \leq j \leq n)$$

and

$$\underline{k}_m \underline{i} \text{ proves } \underline{i} \quad (m \geq 2).$$

Therefore, each α_j ($1 \leq j \leq n$) is provable from $\underline{g}_* := \underline{g}_{(m,n)}^\pi$ by (MP) and (SB) only, and this completes the proof of the Theorem with $r \geq 0$. \square

Unlike in Theorem 12, the hypotheses of Theorem 19 allow also applications to (purely implicative) relevant logics (in the sense of [1], [22], [25], [26], [28]). But, as earlier, in section 3, where we paused on C.A. Meredith's $BCK_{\wedge} \rightarrow$, we prefer to reach that point via some intermediary landmark. The motivation behind this détour will appear soon.

Let $BCK_{\wedge} \rightarrow$ be the Jaśkowski-Meredith purely implicative logic (cf. [18], Appendix I, [14], section 7, or even [17], whose "Postscript" gives the reason we had to use the name above). This is, by the way, a relevant logic in the sense of [1], [22]. Specifically, it coincides with the purely implicative fragment of what the defenders of relevance use to call "Relevance without Contraction" (" $R_{\wedge} \rightarrow$ ", for short, where both " \rightarrow " and "Contraction" denote our formula \underline{w} , that is: "the Hilbert formula" of the post-war Dublin residents, whether Polish or not) and has been studied — on different grounds — by various persons among which S. Jaśkowski, C.A. Meredith (as principal proponents; see references given earlier), A. Church, N.D. Belnap Jr., A. Urquhart (cf. [25], [27]), R. Routley, R.K. Meyer (see [22] and the references given there), and the author ([19]).

By definition, $BCK_{\wedge} \rightarrow$ is finitely axiomatizable in (MP) with, as axioms, \underline{b} , \underline{c} and \underline{i} . C.A. Meredith has also established its Tarski axiomatizability in (MP) (cf. [18], Appendix I, [14], section 7; for alternative axiomatizations see also [19]).

We will be interested in extensions of $BCK_{\wedge} \rightarrow$ still possessing this property.

One has the following straightforward consequence of Theorem 19.

COROLLARY 20.

Any finitely axiomatizable extension L of $BCK_{\wedge} \rightarrow$ in some set of rules $\{(R_1), \dots, (R_r), (MP)\}$, ($r \geq 0$), such that L has a sequentially m -solvable basis, for some $m \geq 2$, is Tarski axiomatizable in the same set of rules.

Proof. Note that

$\underline{c}\underline{i}$ proves $\underline{k}_0 := \underline{c}*$,
by Lemma 5:(5), and, as in Remark 15,

$\underline{b}\underline{c}\underline{c}*$ proves $\underline{d} := \underline{d}_0$

and

$\underline{b}(\underline{b}\underline{c})\underline{d}$ proves \underline{t} .

Also, for all $n \geq 0$,

$\underline{c}(\underline{k}_n \underline{i})$ proves \underline{k}_{n+1} ,

so $\underline{d}, \underline{t}$ and the \underline{k}_m 's ($m \geq 0$) are all theorems of $\text{BCI}_{\lambda \lambda \rightarrow}$.

Finally, apply Theorem 19 to the case in point. \square

To get the Tarski axiomatizability of Church's weak implication (and to show that there are logics satisfying the hypotheses — indeed, somewhat involved — of Corollary 20) recall first that, in [3], $\text{R}_{\lambda \rightarrow}$ was finitely axiomatizable in (MP) with, as axioms, $\underline{w}, \underline{b}, \underline{c}$ and \underline{i} . Henceforth, $\text{R}_{\lambda \rightarrow}$ will denote this formulation of Church's system. (But note that the basis $\{\underline{w}, \underline{b}', \underline{c}, \underline{i}\}$ has the same effect as Church's, in view of Lemma 5: (1) and (2), and similarly, with \underline{c} replaced by either \underline{c}_* or \underline{c}' and/or \underline{w} replaced by \underline{s} or \underline{s} , etc.; see [19] for details.)

Clearly, Church's basis is unsolvable due to the presence of \underline{w} (for \underline{w}_i is already undefined; and similarly, for the remaining four-element bases suggested earlier). So even if $\text{BCI}_{\lambda \lambda \rightarrow}$ is trivially a subsystem of $\text{R}_{\lambda \rightarrow}$, we have no means to apply our Corollary 20 (or, equivalently, Theorem 19) to Church's system unless we can find a solvable basis for it such that it is sequentially so, too. In order to do this and to shorten both the underlying verifications (for sequential solvability) and the length of the resulting single axiom we shall do first of all some axiom chopping.

(The combinatory argument behind half of the following Lemma — its "hard" part — relies on a similar construction which could have been traced back to the work of the pioneers of combinatory logic, viz. to F.B.Fitch's Yale dissertation, 1934.)

Define some more bold face types, namely,

$\underline{a} := \text{CCpCqrCCspCqCsr},$

$\underline{w}' := \text{CpCCpCpqq}.$

Now we have a useful

LEMMA 21.

$\text{R}_{\lambda \rightarrow}$ is finitely axiomatizable in (MP) with, as bases,

- | | |
|--|---|
| (i) $\{\underline{w}, \underline{i}, \underline{a}\}$ | (ii) $\{\underline{w}', \underline{i}, \underline{a}\}$ |
| (iii) $\{\underline{w}', \underline{b}, \underline{c}, \underline{i}\}$ | (iv) $\{\underline{w}', \underline{b}', \underline{c}, \underline{i}\}$ |
| (v) $\{\underline{w}', \underline{b}, \underline{c}_*, \underline{i}\}$ | (vi) $\{\underline{w}', \underline{b}', \underline{c}_*, \underline{i}\}$ |
| (vii) $\{\underline{w}', \underline{b}, \underline{c}', \underline{i}\}$ | (viii) $\{\underline{w}', \underline{b}', \underline{c}', \underline{i}\},$ |
- etc.

Proof. Let, for convenience, $B_i(R_{\lambda \rightarrow})$, with $1 \leq i \leq 8$, be the corresponding bases and $B_0(R_{\lambda \rightarrow})$ be Church's basis $\{\underline{w}, \underline{b}, \underline{c}, \underline{i}\}$.

(i): $B_0(R_{\lambda \rightarrow})$ is contained in $B_1(R_{\lambda \rightarrow})$ for one has that

$$\begin{aligned} \underline{a}\underline{i} & \text{ proves } \underline{c}, \\ \underline{c}\underline{i} & \text{ proves } \underline{c}_*, \\ \underline{a}\underline{c}_* & \text{ proves } \underline{b}', \end{aligned} \quad \text{Lemma 5:(5)}$$

and

$$\underline{c}\underline{b}' \text{ proves } \underline{b}, \quad \text{Lemma 5:(1).}$$

Conversely,

$$\underline{b}(\underline{b}\underline{c})\underline{b} \text{ proves } \underline{a},$$

so $B_0(R_{\lambda \rightarrow})$ contains $B_1(R_{\lambda \rightarrow})$.

(ii): $B_1(R_{\lambda \rightarrow})$ and $B_2(R_{\lambda \rightarrow})$ are equivalent, for

$$\underline{c}\underline{w} \text{ proves } \underline{w}'$$

and

$$\underline{c}\underline{w}' \text{ proves } \underline{w}.$$

(iii)-(viii): Use (ii) and Lemma 5. \square

LEMMA 22.

For all $\alpha, \beta_1, \beta_2, \beta_3, \beta_4$ in any implicative logic L ,

$$(1) \quad \underline{w}'\alpha = \underline{d}\alpha\alpha = \underline{d}_2(\alpha, \alpha);$$

$$(2) \quad \underline{w}'\underline{i}\underline{i} = \underline{i} \text{ (i.e., } \underline{w}' \text{ is 2-solvable or } m\text{-solvable with } m \geq 2);$$

$$(3) \quad \underline{w}\underline{b}\underline{i} = \underline{b}\underline{i}\underline{i} = \underline{i}; \quad (4) \quad \underline{w}\underline{b}'\underline{i} = \underline{b}'\underline{i}\underline{i} = \underline{i};$$

(so \underline{b} and \underline{b}' are 2-solvable and therefore m -solvable for $m \geq 2$).

$$(5) \quad \underline{w}\underline{a}\underline{i} = \underline{a}\underline{i}\underline{i} = \underline{c}_*; \quad (6) \quad \underline{c}_*\underline{i}\underline{i} = \underline{i} \text{ (} \underline{c}_* \text{ is 2-solvable);}$$

$$(7) \quad \underline{a}\beta_1\beta_2\beta_3\beta_4 \sim \beta_1(\beta_2\beta_4)\beta_3 \text{ and } \underline{a} \text{ is } m\text{-solvable with } m \geq 4.$$

$$(8) \quad \underline{c}, \underline{c}' \text{ are 3-solvable (and hence } m\text{-solvable with } m \geq 3).$$

Proof. Trivial. \square

COROLLARY 23.

$$B_2(R_{\lambda \rightarrow}) = \{\underline{w}', \underline{i}, \underline{a}\} \text{ is sequentially 4-solvable.}$$

Proof. As \underline{i} is 0-solvable and, by Lemma 22, \underline{w}' is 2-solvable and \underline{a} is 4-solvable the set $B_2(R_{\lambda \rightarrow})$ is, altogether, m -solvable with $m \geq 4$.

Now set $\underline{h}_1 := \underline{w}', \underline{h}_2 := \underline{d}_2(\underline{w}', \underline{i}) := \underline{c}\underline{c}\underline{w}'\underline{c}\underline{i}\underline{p}_1\underline{p}_1$ and

$\underline{h}_3 := \underline{d}_3(\underline{w}', \underline{i}, \underline{a}) := \underline{c}\underline{c}\underline{c}\underline{c}\underline{w}'\underline{c}\underline{i}\underline{p}_1\underline{p}_1\underline{c}\underline{a}\underline{p}_2\underline{p}_2$, with notational conventions as in the proof of Theorem 19, for the sequence $\underline{w}', \underline{i}, \underline{a}$ (π is the identical permutation on the set $\{1, 2, 3\}$).

Clearly, \underline{h}_1 is 2-solvable, by Lemma 22:(2). Further, one has that

$$\underline{h}_2\underline{i}\underline{i} = \underline{w}'\underline{i}\underline{i} = \underline{i} \quad \text{by Lemma 22:(2),}$$

so \underline{h}_2 is 2-solvable and, finally,

$\underline{h}_3 \underline{i} \underline{i} \underline{i} = \underline{h}_2 \underline{a} \underline{i} \underline{i} = \underline{a} \underline{w}' \underline{i} \underline{i} \underline{i} = \underline{w}' (\underline{i} \underline{i}) \underline{i} = \underline{w}' \underline{i} \underline{i} = \underline{i}$,
by Lemma 1:(11), Remark 3 and Lemma 22:(2),(7), with Corollary 8. Thus \underline{h}_3 is 3-solvable and, altogether, the set $\{\underline{h}_1, \underline{h}_2, \underline{h}_3\}$ is m -solvable with $m \geq 3$. This completes the proof. \square

Now we get the envisaged result, viz.,

COROLLARY 24.

R_{λ} is Tarski axiomatizable in (MP).

Proof. $\underline{B} \underline{C} \underline{I}_{\lambda}$ is a subsystem of R_{λ} (cf. Church's basis) and, by Corollary 23, it has a sequentially 4-solvable basis, namely $\underline{B}_2(R_{\lambda}) = \{\underline{w}', \underline{i}, \underline{a}\}$. So the result follows by Corollary 20.

Explicitely, where \underline{h}_3 is as in the proof of Corollary 23, the single axiom obtained on the pattern of proof of Theorem 19 is

$$\underline{r} := \underline{t}(\underline{k}_4, \underline{h}_3, \underline{k}_4) := \underline{C} \underline{C} \underline{k}_4 \underline{C} \underline{h}_3 \underline{C} \underline{k}_4 \underline{p} \underline{p},$$

as expected. \square

REMARK 25.

$\underline{B}_3(R_{\lambda}) = \{\underline{w}', \underline{i}, \underline{b}, \underline{c}\}$ in Lemma 21 is sequentially 3-solvable, so we might have been obtained an alternative (longer) single axiom starting from this basis of R_{λ} . Indeed, set $\underline{h}_1^* := \underline{w}'$, $\underline{h}_2^* := \underline{d}_2(\underline{w}', \underline{i})$, $\underline{h}_3^* := \underline{d}_3(\underline{w}', \underline{i}, \underline{b})$ and $\underline{h}_4^* := \underline{d}_4(\underline{w}', \underline{i}, \underline{b}, \underline{c})$. It is easy to see that $\underline{h}_1^*, \underline{h}_2^*$ are 2-solvable (cf. the proof of Corollary 23), while \underline{h}_3^* is 3-solvable ($\underline{h}_3 \underline{i}^3 = \underline{h}_2 \underline{b} \underline{i}^2 = \underline{b} \underline{w}' \underline{i}^3 = \underline{w}' \underline{i} \underline{i} = \underline{i}$) and \underline{h}_4^* is 2-solvable (by $\underline{h}_4 \underline{i}^2 = \underline{h}_3^* \underline{c} \underline{i} = \underline{c} \underline{h}_2^* \underline{b} \underline{i} = \underline{h}_2^* \underline{i} \underline{b} = \underline{w}' \underline{i} \underline{b} = \underline{d} \underline{i} \underline{i} \underline{b} = \underline{b} \underline{i} \underline{i} = \underline{i}$). So the new single axiom for R_{λ} would be

$$\underline{r}^* := \underline{t}(\underline{k}_3, \underline{h}_4^*, \underline{k}_3) := \underline{C} \underline{C} \underline{k}_3 \underline{C} \underline{h}_4^* \underline{C} \underline{k}_3 \underline{p} \underline{p}.$$

More variations on this theme are certainly possible.

A shortcoming of Theorem 19 (resp. Corollary 20) consists of that it can be applied directly only to (purely implicative) logics possessing at least one solvable basis. This motivates the following generalization of Theorem 19.

THEOREM 26.

Theorem 19 holds with "sequentially m -solvable" replaced by "sequentially m -quasi-solvable".

Proof. Let L_{λ} be finitely axiomatizable in some set of rules containing (MP), for the specified notion of implication, with axioms

$$\alpha_1, \dots, \alpha_p, \beta_1, \dots, \beta_q \quad (p, q \geq 0, p+q \geq 1),$$

where the α_i 's are unsolvable and the β_j 's are m -solvable, for some $m \geq 2$. Construct now, for $\gamma_1, \dots, \gamma_p$ m -solvable theorems of L_{λ} , formulae

$$\begin{aligned}\underline{v}_i &:= d_2(\alpha_i, \gamma_i) & (1 \leq i \leq p), \\ \underline{f}_i &:= d_i(\underline{v}_1, \dots, \underline{v}_i) & (1 \leq i \leq p), \\ \underline{h}_j &:= d_{j+1}(\underline{f}_p, \beta_1, \dots, \beta_j) & (1 \leq j \leq q).\end{aligned}$$

By the hypothesis of the Theorem these formulae must be m -solvable, too. The needed single axiom for L_λ is now

$\underline{g} := \underline{g}(p, q, m) := t(k_m, h_q, k_m) := C C k_m C h_q C k_m s s$
and one has, as earlier in the proof of Theorem 19, that

$$\underline{g} \underline{g}(\underline{g} \underline{g}) \text{ proves } k_m,$$

for \underline{h}_q is m -solvable (and so is k_m) and

$$\underline{g}(\underline{g} \underline{g}) \text{ proves } h_q.$$

"Sequentially" points out, as in the case of sequential solvability, to the way of obtaining the α_i 's and the β_j 's ($1 \leq i \leq p, 1 \leq j \leq q$).

Specifically, the \underline{h}_j 's ($1 \leq j < q$) and \underline{f}_p can be obtained from \underline{h}_q , i.e.

$$\underline{h}_j k_m \text{ proves } \underline{h}_{j-1}, h_1 k_m \text{ proves } \underline{f}_p \quad (1 \leq j < q),$$

for the β_j 's ($1 \leq j \leq q$) are m -solvable.

Then the \underline{f}_i 's ($1 \leq i < p$) can be obtained from \underline{f}_p as follows:

$$\underline{f}_i k_m \text{ proves } \underline{f}_{i-1} \quad (1 < i \leq p),$$

for the \underline{v}_i 's ($1 < i \leq p$) are m -solvable. (Note that $\underline{f}_1 := \underline{v}_1$.)

Now the remaining \underline{v}_i 's ($1 < i \leq p$) can be obtained from the \underline{f}_i 's, i.e.

$$\underline{f}_i (k_m k_m) k_m \text{ proves } \underline{v}_i \quad (1 < i \leq p),$$

for the \underline{f}_i 's ($1 \leq i < p$) are m -solvable.

Finally, the α_i 's can be obtained from the \underline{v}_i 's ($1 \leq i \leq p$), since the γ_i 's are supposed to be m -solvable and hence

$$\underline{v}_i k_m \text{ proves } \alpha_i \quad (1 \leq i \leq p),$$

while the β_j 's ($1 \leq j \leq q$) can be obtained from the corresponding \underline{h}_j 's, as expected, i.e., by

$$\underline{h}_j (k_m k_m) k_m \text{ proves } \beta_j \quad (1 \leq j \leq q),$$

for $\underline{h}_1, \dots, \underline{h}_{q-1}$ are m -solvable.

On the other hand, $\underline{g} := \underline{g}(p, q, m)$ is a theorem of L_λ , for so is \underline{h}_q (the γ_i 's, $1 \leq i \leq p$, were supposed to be theorems of L_λ , while \underline{g} is a theorem of L_λ , by the hypothesis of the Theorem), k_m and t . This completes the proof (Lemmas 10 and 11 were used tacitly). \square

REMARK 27.

An analogue of Corollary 20 holds with "sequentially m -solvable" replaced by "sequentially m -quasi-solvable".

REMARK 28.

As an application of Theorem 26 (resp. Corollary 27) consider Church's axiomatization of $R_{\lambda \rightarrow}$ with

$$\alpha_1 := \underline{w}, \beta_1 := \underline{i}, \beta_2 := \underline{c}, \beta_3 := \underline{b}.$$

Set $\gamma_1 := \underline{b}'$ (recall that $\underline{c}\underline{b}$ proves \underline{b}').

Clearly, $\underline{v}_1 := \underline{f}_1 := \underline{d}_2(\underline{w}, \underline{b}')$. Now, with the conventions of the proof of Theorem 26, one has

$$\begin{aligned} \underline{h}_1 &:= \underline{d}_2(\underline{f}_1, \underline{i}) := \underline{d}_3(\underline{w}, \underline{b}', \underline{i}), \\ \underline{h}_2 &:= \underline{d}_3(\underline{f}_1, \underline{i}, \underline{c}) := \underline{d}_4(\underline{w}, \underline{b}', \underline{i}, \underline{c}), \end{aligned}$$

and

$$\underline{h}_3 := \underline{d}_4(\underline{f}_1, \underline{i}, \underline{c}, \underline{b}) := \underline{d}_5(\underline{w}, \underline{b}', \underline{i}, \underline{c}, \underline{b}).$$

Using now Lemma 4, Remark 3, Corollary 8 and Lemma 22:(4) one can show that $\underline{f}_1, \underline{h}_1$ and \underline{h}_2 are 2-solvable, while \underline{h}_3 is 4-solvable. As the set of β_j 's ($j = 1, 2, 3$) is 3-solvable and $\gamma_1 := \underline{b}'$ is 2-solvable (cf. Lemma 22:(3), (8), etc.), we have already established that Church's basis is sequentially 4-quasi-solvable. On the other hand, we know that $\underline{d}, \underline{t}$, and \underline{k}_4 are $R_{\lambda \rightarrow}$ -theorems (cf. the proof of Corollary 20), so Church's system is — again — Tarski axiomatizable in (MP) with, as single axiom,

$$\underline{r}_{\&} := \underline{t}(\underline{k}_4, \underline{h}_3, \underline{k}_4) := \underline{CCK}_4 \underline{Ch}_3 \underline{CK}_4 \underline{pp},$$

where \underline{h}_3 is as in this Remark.

REMARK 29.

It is possible to shorten the latter single axiom $\underline{r}_{\&}$, obtained in Remark 28, using the fact that $\underline{a}\underline{i}\underline{i}$ proves \underline{c}_* (cf. Lemma 22:(5)). Indeed, set, with the conventions of the proof of Theorem 26,

$$\alpha_1 := \underline{w}, \beta_1 := \underline{i}, \beta_2 := \underline{b} \text{ and } \gamma_1 := \underline{a}.$$

(We have seen in Lemma 21 that \underline{a} is a theorem of $\text{BCI}_{\lambda \rightarrow}$ and hence of $R_{\lambda \rightarrow}$.)

Now construct formulae $\underline{v}_1 := \underline{d}_2(\underline{w}, \underline{a}), \underline{f}_1 := \underline{v}_1, \underline{h}_1 := \underline{d}_2(\underline{f}_1, \underline{i}) := \underline{d}_3(\underline{w}, \underline{a}, \underline{i})$ and $\underline{h}_2 := \underline{d}_3(\underline{f}_1, \underline{i}, \underline{b}) := \underline{d}_4(\underline{w}, \underline{a}, \underline{i}, \underline{b})$. One finds easily that $\underline{h}_2 \underline{i}^3$ proves \underline{c}_* (for $\underline{h}_2 \underline{i}^3 = \underline{h}_1 \underline{b} \underline{i} \underline{i} = \underline{b} \underline{f}_1 \underline{i} \underline{i} \underline{i} = \underline{f}_1 (\underline{i} \underline{i}) \underline{i} = \underline{f}_1 \underline{i} \underline{i} = \underline{w} \underline{a} \underline{i} = \underline{a} \underline{i} \underline{i} = \underline{c}_*$); so one has also that $\underline{h}_2 \underline{i}^5 = \underline{i}$, for \underline{c}_* is 2-solvable. Next \underline{h}_1 is 4-solvable for $\underline{h}_1 \underline{i}^4 = \underline{f}_1 \underline{i}^4 = \underline{w} \underline{a} \underline{i}^3 = \underline{a} \underline{i}^4 = \underline{i}$ and, finally, $\underline{v}_1 := \underline{f}_1$ is 4-solvable, by the same token ($\underline{f}_1 \underline{i}^4 = \underline{w} \underline{a} \underline{i}^3$, etc.). So the set $\{\underline{w}, \underline{i}, \underline{b}\}$ is sequentially 5-quasi-solvable and we may set as single axiom (for $R_{\lambda \rightarrow}$) $\underline{r}_{\&} := \underline{CCK}_5 \underline{Ch}_2 \underline{CK}_5 \underline{pp}$, with \underline{h}_2 as above. The additional trick (beyond the pattern of proof of Theorem 26) consists of getting \underline{c}_* from \underline{h}_2 and \underline{i} . But the set $\{\underline{w}, \underline{i}, \underline{b}, \underline{c}_*\}$ is a basis for $R_{\lambda \rightarrow}$ (by Lemma 5).

Theorem 19 admits now of a generalization in some other direction, viz. by generalizing the concept of solvability.

Let L_\wedge be some (purely) implicative logic and $\vec{\beta} := \beta_1, \dots, \beta_m$ be a sequence of theorems of L_\wedge (possibly with repetitions and possibly empty). For $\vec{\beta}$ fixed set

$$k_{\vec{\beta}} := CpCC\beta_1 \dots C\beta_m CCpqq.$$

(One can see easily that for $\vec{\beta}$ empty one has $k_{\vec{\beta}} := k_0 := c_*$.)

A set $\{\alpha_1, \dots, \alpha_n\}$ of theorems of L_\wedge is $\vec{\beta}$ -solvable (for $\vec{\beta}$ fixed) if $\alpha_j \vec{\beta} := \alpha_j \beta_1 \dots \beta_m$ proves \underline{i} for all $j, 1 \leq j \leq n$.

Construct now for some sequence $\alpha_1, \dots, \alpha_n$ (of theorems of L_\wedge) formulae $\underline{h}_j := d_j(\alpha_1, \dots, \alpha_j)$ with $1 \leq j \leq n$.

A set $\{\alpha_1, \dots, \alpha_n\}$ of theorems of L_\wedge is sequentially $\vec{\beta}$ -solvable (for $\vec{\beta}$ fixed) if its elements can be arranged (without repetitions) in a sequence $\alpha_1, \dots, \alpha_n$ say such that

- (i) each α_j is $\vec{\beta}$ -solvable ($1 \leq j \leq n$),
- (ii) each \underline{h}_j (constructed as above) is $\vec{\beta}$ -solvable ($1 \leq j \leq n$)

and, finally,

- (iii) $k_{\vec{\beta}}$ is $\vec{\beta}$ -solvable.

Clearly, if each β_i in $\vec{\beta}$ ($1 \leq i \leq m$) is the formula \underline{i} , $\vec{\beta}$ -solvability for this $\vec{\beta}$ amounts to m -solvability and sequential $\vec{\beta}$ -solvability coincides with sequential m -solvability for $m \geq 2$.

One can prove by a straightforward extension of the methods used earlier that the following generalization of Theorem 19 holds.

THEOREM 30.

Theorem 19 holds with "sequential m -solvability" replaced by "sequential $\vec{\beta}$ -solvability" for some fixed sequence $\vec{\beta} := \beta_1, \dots, \beta_m$ of theorems of L_\wedge and with " k_m is a theorem of L_\wedge " replaced by " $k_{\vec{\beta}}$ is a theorem of L_\wedge ".

Proof. Mutatis mutandis, as for Theorem 19. \square

REMARK 31.

Note also that Theorem 19 becomes a particular case of Theorem 30 with $\vec{\beta} := \underline{i}_1, \dots, \underline{i}_m$ (the \underline{i}_j 's are lexical variants of \underline{i} , as in section 1 above).

Theorems 19, 26 and 30 do not apply as such to the purely implicative fragment E_{\rightarrow} of the Anderson-Belnap Entailment system (cf. [1]) due to the fact $\underline{d}, \underline{t}$ and the \underline{k}_m 's ($m \geq 0$) are not theorems of E_{\rightarrow} . But the pattern of proof used earlier still works for some slight modification of the corresponding hypotheses.

Define first, with $\hat{p} := Cp'p''$, $\hat{q} := Cq'q''$ and $\hat{r} := Cr'r''$, the following formulae:

$$\begin{aligned}\hat{\underline{d}} &:= C\hat{p}C\hat{q}CC\hat{p}C\hat{q}rr, \\ \hat{\underline{t}} &:= C\hat{p}C\hat{q}C\hat{r}CC\hat{p}C\hat{q}C\hat{r}ss, \\ \hat{\underline{k}}_0 &:= \hat{\underline{c}}_* := C\hat{p}CC\hat{p}qq, \\ \hat{\underline{k}}_n &:= C\hat{p}CC\underline{c}_1 \dots \underline{c}_n CC\hat{p}qq, & (n \geq 1), \\ \hat{\underline{c}} &:= CCpCqrCqCpr.\end{aligned}$$

(All these formulae are theorems of E_{\rightarrow} , as one can easily check using the Fitch style formulation of E_{\rightarrow} in [1]. But see Corollary 33 below for the corresponding Hilbert style derivations, with condensed detachment.)

We can now establish the following (stronger) form of Theorem 19.

THEOREM 32.

Theorem 19 holds with \underline{k}_m ($m \geq 2$), $\underline{d}, \underline{t}$ replaced by $\hat{\underline{k}}_m, \hat{\underline{d}}, \hat{\underline{t}}$ resp.

Proof. As earlier, for Theorem 19. \square

Let now $B_{\rightarrow}^{\hat{\underline{c}}}$ be the (purely) implicative logic axiomatized by $\underline{b}, \hat{\underline{c}}$ and \underline{i} with (MP) and (SB) as primitive rules of derivation. It is known that $B_{\rightarrow}^{\hat{\underline{c}}}$ can be axiomatized also with (MP), (SB) and, as axioms, \underline{b}' and $\underline{q} := \underline{k}_0$, i.e.,

$$\underline{q} := CCCppqq,$$

(cf., e.g., [14] or [1], 8.5.1.; the result is due to M. Wajsberg, C.A. Meredith and N. Belnap Jr., independently). C.A. Meredith has also found that $B_{\rightarrow}^{\hat{\underline{c}}}$ is Tarski axiomatizable in (MP) (cf. [14], section 10 or [1], 8.5.1.).

It can be easily seen that the following (stronger) form of Corollary 20 obtains.

COROLLARY 33.

Corollary 20 holds with BCI_{\rightarrow} replaced by $B_{\rightarrow}^{\hat{\underline{c}}}$.

Proof. Note that $\underline{b}\hat{\underline{c}}\hat{\underline{c}}_*$ proves $\hat{\underline{d}}, \underline{b}(\underline{b}\hat{\underline{c}})\hat{\underline{d}}$ proves $\hat{\underline{t}}, \hat{\underline{k}}_0 := \hat{\underline{c}}_*$ and $\hat{\underline{c}}\underline{i}$ proves $\hat{\underline{c}}_*$, while, for all $m \geq 0$, $\hat{\underline{c}}(\hat{\underline{k}}_m \underline{i})$ proves $\hat{\underline{k}}_{m+1}$. Then apply Theorem 32. \square

One has also the following (stronger) analogues of Theorem 26 and Remark 27.

THEOREM 34.

Theorem 19 holds with $\underline{d}, \underline{t}, \underline{k}_m$ replaced by $\hat{\underline{d}}, \hat{\underline{t}}, \hat{\underline{k}}_m$ resp. ($m \geq 2$) and "sequentially m -solvable" replaced by "sequentially m -quasi-solvable".

Proof. Mutatis mutandis, as for Theorem 26. \square

COROLLARY 35.

Corollary 20 holds with $\underline{BCI}_{\lambda \rightarrow}$ replaced by $\hat{\underline{BCI}}_{\lambda \rightarrow}$ and "sequentially m -solvable" replaced by "sequentially m -quasi-solvable".

Proof. Note that $\hat{\underline{d}}, \hat{\underline{t}}$, and the $\hat{\underline{k}}_m$'s ($m \geq 0$) are theorems of $\hat{\underline{BCI}}_{\lambda \rightarrow}$, then apply Theorem 34. \square

Recall now (from [1], say) that the following sets of formulae axiomatize the Pure Entailment system $\underline{E}_{\lambda \rightarrow}$ with (MP) and (SB) as primitive rules of derivation:

$$\underline{B}_0(\underline{E}_{\lambda \rightarrow}) = \{\underline{w}, \underline{b}, \hat{\underline{c}}, \underline{i}\}, \quad \underline{B}_1(\underline{E}_{\lambda \rightarrow}) = \{\underline{w}, \underline{b}', \underline{o}\}$$

(the latter one is Belnap's preferred basis for $\underline{E}_{\lambda \rightarrow}$). Clearly, $\underline{E}_{\lambda \rightarrow}$ is a (proper) extension of $\hat{\underline{BCI}}_{\lambda \rightarrow}$ (though not of $\underline{BCI}_{\lambda \rightarrow}$) and we may readily apply Corollary 35 to Belnap's basis $\underline{B}_1(\underline{E}_{\lambda \rightarrow})$ say, getting the expected result, viz.,

COROLLARY 36.

$\underline{E}_{\lambda \rightarrow}$ is Tarski axiomatizable in (MP).

Proof. (Nearly completed above. Still, for the sake of effectiveness we can afford the following considerations.)

Set $\alpha_1 := \underline{w}, \beta_1 := \underline{o}, \beta_2 := \underline{b}'$ and $\gamma_1 := \underline{b}'$. Further, with conventions as in the proof of Theorem 26 (resp. Theorem 34), set $\underline{v}_1 := \underline{f}_1 := \underline{d}_2(\underline{w}, \underline{b}'), \underline{h}_1 := \underline{d}_2(\underline{f}_1, \underline{o}) := \underline{d}_3(\underline{w}, \underline{b}', \underline{o})$ and $\underline{h}_2 := \underline{d}_3(\underline{f}_1, \underline{o}, \underline{b}') := \underline{d}_4(\underline{w}, \underline{b}', \underline{o}, \underline{b}')$. Now \underline{h}_2 is 2-solvable for $\underline{h}_2 \underline{i} \underline{i} = \underline{h}_1 \underline{b}' \underline{i} = \underline{b}' \underline{f}_1 \underline{o} \underline{i} = \underline{o}(\underline{f}_1 \underline{i}) = \underline{o}(\underline{w} \underline{b}') = \underline{w} \underline{b}' \underline{i} = \underline{b}' \underline{i} \underline{i} = \underline{i}$; \underline{h}_1 is 1-solvable for $\underline{h}_1 \underline{i} = \underline{f}_1 \underline{o} = \underline{o}(\underline{w} \underline{b}') = \underline{w} \underline{b}' \underline{i}$, etc. and $\underline{v}_1 := \underline{f}_1$ is 2-solvable ($\underline{f}_1 \underline{i} \underline{i} = \underline{w} \underline{b}' \underline{i}$, etc.). Finally, \underline{b}' is 2-solvable and \underline{o} is 1-solvable. Thus Belnap's basis is sequentially 2-quasi-solvable and the single axiom for $\underline{E}_{\lambda \rightarrow}$ might be now, with \underline{h}_2 as above,

$$\underline{e} := \underline{t}(\underline{k}_2, \underline{h}_2, \underline{k}_2) := \underline{CCK}_2 \underline{Ch}_2 \underline{CK}_2 \underline{pp},$$

on the usual pattern employed earlier for $\underline{R}_{\lambda \rightarrow}$. \square

REMARK 37.

As expected, a (stronger) variant of Theorem 30 holds with $\hat{k}_{\vec{\beta}} := CCp'p'CC\beta_1 \dots C\beta_m CCCp'p'qq$ instead of $\hat{k}_{\vec{\beta}}$, for some fixed sequence $\vec{\beta} := \beta_1, \dots, \beta_m$ ($m \geq 2$ say) of formulae of $L_{\hat{\lambda}}$. The corresponding statement is an obvious generalization of Theorem 32.

REMARK 38.

Unlike the methods of proof available in the presence of \hat{k} (in section 3 above) the methods of finding single axioms used in this section do not apparently apply to implicative logics containing conjunction and/or disjunction (these ingredients would obviously block the application of the unification algorithm while evaluating pe's). In particular, the full Relevance logic $R_{\hat{\lambda}}$ (cf. [1], [22], [25]), the Entailment system $E_{\hat{\lambda}}$ of Anderson-Belnap (cf. [1], [25]) and the Prawitz-Urquhart system $S_{\hat{\lambda}}$ (cf. [28], say) as well as many other (propositional) relevant logics reviewed in [1] and [22] are cases in point. Specifically, it is an open problem whether the relevant logics $R_{\hat{\lambda}}, E_{\hat{\lambda}}, S_{\hat{\lambda}}$, etc. are Tarski axiomatizable in (MP) and the Adjunction rule (ADJ):

$$\alpha, \beta \implies K\alpha\beta$$

(where K is Polish notation for conjunction).

(It should be however obvious how to obtain 2-axioms bases for these logics, with (MP), (ADJ) and (SB) as primitive rules, due to the presence of CKpqp and/or CKpqq.)

But note that the CN-fragments (i.e., those containing implication and negation only) of the relevant logics above (as well as many other similar systems not named here) are Tarski axiomatizable in (MP).

31 December 1981

Alex. van Humboldtstraat 4,
3514 GP Utrecht,
The Netherlands.

REFERENCES

ANDERSON, A.R., BELNAP JR., N.D. et alii

- [1] Entailment. The Logic of Relevance and Necessity, Volume I,
Princeton University Press; Princeton, 1975.

BARENDREGT, H.P.

- [2] The Lambda Calculus, Its Syntax and Semantics,
North Holland; Amsterdam, New York, Oxford (Studies in Logic and the Foundations of Mathematics 103), 1981

CHURCH, A.

- [3] The theory of weak implication,
in MENNE, A., WILHEMY, A. and ANGSI, H. (eds.) Kontrolliertes Denken, Untersuchungen zum Logikkalkül und zur Logik der Einzelwissenschaften, Kommissionsverlag Karl Alber; Munich, 1951, pp. 22-37.

CURRY, H.B., FEYS, R. and CRAIG, W.

- [4] Combinatory Logic, Volume I,
North Holland; Amsterdam, 1958 (reprinted 1974).

VAN DALEN, D.

- [5] Intuitionistic logic,
in GABBAY, D.M. and GUENTHNER, F. (eds.) Philosophical Logic, D. Reidel; Dordrecht [forthcoming].
(First draft available as: Preprint nr. 209, University of Utrecht, Department of Mathematics, September 1981.)

HINDLEY, J.R., LERCHER, B. and SELDIN, J.P.

- [6] Introduction to Combinatory Logic,
Cambridge University Press; London (London Math. Soc. Lecture Notes 7), 1972.

HOWARD, W.

- [7] The Formulae-as-Types notion of construction,
in SELDIN, J.P. and HINDLEY, J.R. (eds.) To H.B. Curry: Essays on Combinatory Logic, Lambda Calculus and Formalism, Academic Press; London, New York, etc., 1980, pp. 479-490
(written in 1969).

KALMAN, J.A.

- [8] An algorithm for Meredith's condensed detachment (abstract),
J. Symbolic Logic 39, 1974, p. 206.

LEŚNIEWSKI, S.

- [9] Grundzüge eines neuen Systems der Grundlagen der Mathematik,
Fund. Mathematicae 14, 1929, pp. 1-81.

LORENZEN, P.

- [10] Einführung in die operative Logik und Mathematik,
Springer Verlag; Berlin, Göttingen, Heidelberg, 1955.

ŁUKASIEWICZ, J.

- [11] The shortest axiom of the implicational calculus,
Proc. Royal Irish Acad., (Section A) 52, 1948, pp. 25-33.
(Reprinted in: ŁUKASIEWICZ, J. Selected Works, edited by
L. BORKOWSKI, North Holland; Amsterdam and Polish Scientific
Publishers, Warsaw, 1970, pp. 295-305.)

- [12] Elements of Mathematical Logic,
Pergamon Press, Macmillan & Co.; New York 1964.

ŁUKASIEWICZ, J. and TARSKI, A.

- [13] Untersuchungen über den Aussagenkalkül,
Comptes Rendus des Séances de la Société des Sciences
et des Lettres de Varsovie, (Classe III - 3^{ème}) 23, 1930, pp.
39-50.
(Reprinted in: ŁUKASIEWICZ, J. Selected Works, edited by
L. BORKOWSKI, North Holland; Amsterdam and Polish Scientific
Publishers, Warsaw, 1970, pp. 131-152 and in TARSKI, A.
Logic, Semantics, Metamathematics, Papers from 1923 to 1938,
Clarendon Press; Oxford, 1956, pp. 38-59.)

MEREDITH, C.A. and PRIOR, A.N.

- [14] Notes on the axiomatics of the propositional calculus,
Notre Dame J. Formal Logic 4, 1963, pp. 171-187.

MEREDITH, D.

- [15] Combinatory and propositional logic,
Notre Dame J. Formal Logic 15, 1974, pp. 156-160.
[16] Carew Arthur Meredith (1904-1970),
Notre Dame J. Formal Logic 18, 1977, pp. 513-516.
[17] Positive logics and λ -constants,
Studia Logica 37, 1978, pp. 269-285.

PRIOR, A.N.

- [18] Formal Logic,
Clarendon Press; Oxford, 1962 (second edition).

REZUS, A.

- [19] Singleton bases for Λ_K^0 and a result of Alfred Tarski,
Preprint nr. 150, University of Utrecht, Department of
Mathematics, April 1980.

ROBINSON, J.A.

- [20] A machine oriented logic based on the resolution principle,
J. Assoc. Comput. Mach. 12, 1965, pp. 23-41

ROBINSON, J.A. (continued)

- [21] Logic: Form and Function. The Mechanization of Deductive Reasoning,

Edinburgh University Press; Edinburgh, 1979.

ROUTLEY, R. and MEYER, R.K.

- [22] Relevant Logics and Their Rivals,

The Australian National University, Research School of Social Sciences Monograph Series (Department of Philosophy); Canberra [forthcoming].

SOBOCIŃSKI, B.

- [23] Z badań nad teorią dedukcji,

Przegląd Filozoficzny 35, 1932, pp. 171-193.

- [24] On well constructed axiom systems,

in Yearbook of the Polish Society of Art and Sciences Abroad, 1955/1956, pp. 54-65.

URQUHART, A.

- [25] The Semantics of Entailment,

Dissertation: University of Pittsburgh, 1973.

- [26] Semantics for relevant logics,

J. Symbolic Logic 37, 1972, pp. 159-169.

- [27] Proofs, snakes and ladders,

Dialogue 13, 1974, pp. 723-731.

- [28] What is relevant implication?

Preprint, University of Toronto, Department of Philosophy, 1974 (with an "Addendum", dated April 1981).

Appendix:

On a Singleton Basis for the Set of Closed Lambda-terms.

As reported in [1, page 180], C.A. Meredith maintained once that the following lambda-term

$$\underline{G} =: \lambda xyz. y(\lambda u. z)(xz)$$

is a basis for the set of closed lambda-(K-)terms. Still he did never supply a proof of this claim in print and several attempts to a reconstruction of the missing derivations (see, e.g., [2, page 283] or [3, pp. 10-11]), even with the help of a computer (programmed in 16K LISP by Professor W.L. van der Poel), have been unsuccessful so far.

Actually, the needed argument is relatively simple.

Recall first some current notation in [3]:

$$\begin{aligned} \underline{I} &=: \lambda x. x, \quad \underline{K} =: \lambda xy. x, \quad \underline{K}' =: \lambda xy. y, \quad \underline{B} =: \lambda xyz. x(yz), \\ \underline{B}' &=: \lambda xyz. y(xz), \quad \underline{C} =: \lambda xyz. xzy, \quad \underline{C}_* =: \lambda xy. yx, \quad \underline{W} =: \lambda xy. xyy, \\ \underline{W}_* &=: \lambda x. xx, \quad \underline{S} =: \lambda xyz. xz(yz), \quad \underline{S}' =: \lambda xyz. yz(xz). \end{aligned}$$

Further, set for any closed lambda-term X , $X_1 =: X$, $X_{n+1} =: X_n X$ (n a positive integer). $=$ will denote beta-convertibility.

Note first that $\underline{G}_3 = \lambda xy. y(xx)$. One has immediately that

$$\underline{C}_* = \underline{G}\underline{G}_3\underline{G}, \quad \underline{I} = \underline{G}\underline{C}_*\underline{C}_*, \quad \underline{K}' = \underline{G}\underline{I}\underline{C}_*\underline{I} \quad \text{and} \quad \underline{K} = \underline{G}\underline{K}'\underline{C}_*.$$

Now one can obtain \underline{B}' , \underline{B} and \underline{C} . Indeed, set $\underline{E} =: \underline{G}(\underline{K}\underline{C}_*)\underline{G}$. (Note that $\underline{E} = \lambda xy. x(Ky) = \underline{B}'\underline{K}$.)

Then $\underline{B}' = \underline{G}_2(\underline{K}\underline{E}) = \underline{E}\underline{G}_2\underline{E}$, $\underline{B} = \underline{B}'\underline{C}_*(\underline{B}'\underline{B}')$ and $\underline{C} = \underline{B}'\underline{B}'(\underline{B}'\underline{C}_*)$.

Finally, one needs any one of \underline{W}_* , \underline{W} , \underline{S}' or \underline{S} . The former two are easy to get: take, e.g., $\underline{W}_* = \underline{G}\underline{I}(\underline{G}_3\underline{I}) = \underline{G}\underline{I}(\underline{C}_*\underline{I})$ or $\underline{W} = \underline{G}_2(\underline{K}(\underline{C}_*\underline{G}_3))$.

For \underline{S}' and \underline{S} one may proceed as in [3], viz. by realizing that

$$\underline{S}' = \underline{B}'\underline{G}(\underline{B}'(\underline{B}'(\underline{C}_*\underline{G}))) \quad \text{and, finally,} \quad \underline{S} = \underline{C}\underline{S}'.$$

Question: is there any basis shorter than \underline{G} ?

REFERENCES

- [1] C.A. MEREDITH, A.N. PRIOR
Notes on the axiomatics of the propositional calculus
 Notre Dame J. Formal Logic 4, 1963, pp. 171-187.
- [2] D. MEREDITH
Positive logic and λ -constants
 Studia Logica 37, 1978, pp. 269-285.
- [3] A. REZUS
Singleton bases for subsets of Λ_K^0 and a result of A. Tarski
 University of Utrecht, Department of Mathematics, Preprint 150,
 April 1980.

Utrecht, 9 October 1981

